

Preprint, arXiv:1010.2489

PROOF OF THREE CONJECTURES ON CONGRUENCES

HAO PAN AND ZHI-WEI SUN

Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China
haopan79@yahoo.com.cn, zwsun@nju.edu.cn

ABSTRACT. In this paper we prove three conjectures on congruences involving central binomial coefficients or Lucas sequences. Let p be an odd prime and let a be a positive integer. We show that if $p \equiv 1 \pmod{4}$ or $a > 1$ then

$$\sum_{k=0}^{\lfloor \frac{3}{4}p^a \rfloor} \binom{-1/2}{k} \equiv \left(\frac{2}{p^a} \right) \pmod{p^2},$$

where $(-)$ denotes the Jacobi symbol. This confirms a conjecture of the second author. We also confirm a conjecture of R. Tauraso by showing that

$$\sum_{k=1}^{p-1} \frac{L_k}{k^2} \equiv 0 \pmod{p} \quad \text{provided } p > 5,$$

where the Lucas numbers $L_0, L_1, L_2, \dots$ are defined by $L_0 = 2$, $L_1 = 1$ and $L_{n+1} = L_n + L_{n-1}$ ($n = 1, 2, 3, \dots$). Our third theorem states that if $p \neq 5$ then we can determine $F_{p^a - (\frac{p^a}{5})} \pmod{p^3}$ in the following way:

$$\sum_{k=0}^{p^a-1} (-1)^k \binom{2k}{k} \equiv \left(\frac{p^a}{5} \right) (1 - 2F_{p^a - (\frac{p^a}{5})}) \pmod{p^3},$$

which appeared as a conjecture in a paper of Sun and Tauraso.

1. INTRODUCTION

In this paper we aim to prove three conjectures on congruences. Our first theorem confirms Conjecture 1.3(i) of Sun [S2].

2010 *Mathematics Subject Classification.* Primary 11B65, 11A07; Secondary 05A10, 11B39.

The second author is the corresponding author. The first author and the second author are respectively supported by the National Natural Science Foundation for Youths in China (grant 10901078), and the National Natural Science Foundation (grant 10871087) and the Overseas Cooperation Fund (grant 10928101) of China.

Theorem 1.1. *Let p be an odd prime and let a be a positive integer. If $p \equiv 1 \pmod{4}$ or $a > 1$, then we have*

$$\sum_{k=0}^{\lfloor \frac{3}{4}p^a \rfloor} \binom{-1/2}{k} \equiv \left(\frac{2}{p^a} \right) \pmod{p^2}, \quad (1.1)$$

where $(-)$ denotes the Jacobi symbol.

Our second theorem confirms a conjecture of R. Tauraso [T].

Theorem 1.2. *Let $p > 5$ be a prime. Then*

$$\sum_{k=1}^{p-1} \frac{L_k}{k^2} \equiv 0 \pmod{p}, \quad (1.2)$$

where the Lucas numbers $L_0, L_1, L_2, \dots$ are defined by $L_0 = 2$, $L_1 = 1$ and $L_{n+1} = L_n + L_{n-1}$ ($n = 1, 2, 3, \dots$).

The Fibonacci sequence $\{F_n\}_{n \geq 0}$ is given by $F_0 = 0$, $F_1 = 1$ and $F_{n+1} = F_n + F_{n-1}$ ($n = 1, 2, 3, \dots$). It is well-known that $p \mid F_{p-(\frac{p}{5})}$ for any odd prime p . Our following theorem determines $F_{p-(\frac{p}{5})} \pmod{p^3}$ and the result appeared as Conjecture 1.1 of Sun and Tauraso [ST].

Theorem 1.3. *Let $p \neq 2, 5$ be a prime and let a be a positive integer. Then we have*

$$\sum_{k=0}^{p^a-1} (-1)^k \binom{2k}{k} \equiv \left(\frac{p^a}{5} \right) (1 - 2F_{p^a-(\frac{p^a}{5})}) \pmod{p^3}. \quad (1.3)$$

Those primes $p > 5$ satisfying the congruence $F_{p-(\frac{p}{5})} \equiv 0 \pmod{p^2}$ are called Wall-Sun-Sun primes (cf. [CDP], [CP, p. 32] and [SS]). Now it is known that there are no Wall-Sun-Sun primes below 2×10^{14} (cf. [MR]).

To understand our proofs of Theorems 1.1-1.3 one needs some basic knowledge of Lucas sequences.

Let A and B be two integers. The Lucas sequence $u_n = u_n(A, B)$ ($n \in \mathbb{N} = \{0, 1, 2, \dots\}$) and its companion $v_n = v_n(A, B)$ ($n \in \mathbb{N}$) are defined by

$$u_0 = 0, \quad u_1 = 1, \quad u_{n+1} = Au_n - Bu_{n-1} \quad \text{for } n = 1, 2, 3, \dots,$$

and

$$v_0 = 2, \quad v_1 = A, \quad v_{n+1} = Av_n - Bv_{n-1} \quad \text{for } n = 1, 2, 3, \dots$$

Let $\Delta = A^2 - 4B$. The characteristic equation $x^2 - Ax + B = 0$ has two roots

$$\alpha = \frac{A + \sqrt{\Delta}}{2} \quad \text{and} \quad \beta = \frac{A - \sqrt{\Delta}}{2},$$

which are both algebraic integers. It is well known that

$$(\alpha - \beta)u_n = \alpha^n - \beta^n \quad \text{and} \quad v_n = \alpha^n + \beta^n \quad \text{for all } n = 0, 1, 2, \dots$$

For an odd prime p and a positive integer a , clearly

$$v_{p^a} \equiv (\alpha + \beta)^{p^a} = A^{p^a} \equiv A \pmod{p}$$

and

$$\Delta u_{p^a} = (\alpha - \beta)(\alpha^{p^a} - \beta^{p^a}) \equiv (\alpha - \beta)^{p^a+1} = \Delta^{(p^a+1)/2} \equiv \Delta \left(\frac{\Delta}{p^a} \right) \pmod{p}.$$

It is also known that $p^a \mid u_{p^a - (\frac{\Delta}{p^a})}$ provided that $p \nmid B$ (see, e.g., [S1, Lemma 2.3]).

Note that $F_n = u_n(1, -1)$ ($n \in \mathbb{N}$) and $L_n = v_n(1, -1)$ ($n \in \mathbb{N}$) are familiar Fibonacci numbers and Lucas numbers respectively. The Pell sequence and its companion are given by $P_n = u_n(2, -1)$ ($n \in \mathbb{N}$) and $Q_n = v_n(2, -1)$ ($n \in \mathbb{N}$) respectively.

We will show Theorems 1.1-1.3 in Sections 2-4 respectively. In the proofs of Theorems 1.2 and 1.3, we employ the useful technique of A. Granville [Gr] and deal with congruences in the ring of algebraic integers.

2. PROOF OF THEOREM 1.1

Lemma 2.1. *Let p be an odd prime and let a be a positive integer. Then*

$$P_{p^a - (\frac{2}{p^a})} Q_{p^a - (\frac{2}{p^a})} \equiv \left(\frac{2}{p^a} \right) \frac{Q_{p^a} - 2}{2} \pmod{p^2}. \quad (2.1)$$

Proof. Recall that $P_n = u_n(2, -1)$ and $Q_n = v_n(2, -1)$ for all $n \in \mathbb{N}$. So

$$P_{p^a} \equiv \left(\frac{2}{p^a} \right) \pmod{p} \quad \text{and} \quad Q_{p^a} \equiv 2 \pmod{p}.$$

Since $Q_{n-1} = 4P_n - Q_n$ and $Q_{n+1} = 4P_n + Q_n$ for $n = 1, 2, 3, \dots$, we have

$$\left(\frac{2}{p^a} \right) Q_{p^a - (\frac{2}{p^a})} = 4 \left(\frac{2}{p^a} \right) P_{p^a} - Q_{p^a} \equiv 2 \pmod{p}.$$

Similarly,

$$P_{p^a - (\frac{2}{p^a})} = \frac{Q_{p^a}}{2} - \left(\frac{2}{p^a}\right) P_{p^a} \equiv 0 \pmod{p}.$$

Clearly $\alpha = 1 + \sqrt{2}$ and $\beta = 1 - \sqrt{2}$ are the two roots of the equation $x^2 - 2x - 1 = 0$. Thus

$$Q_n^2 - 8P_n^2 = (\alpha^n + \beta^n)^2 - (\alpha^n - \beta^n)^2 = 4(\alpha\beta)^n = 4(-1)^n$$

for all $n \in \mathbb{N}$. Therefore

$$Q_{p^a - (\frac{2}{p^a})}^2 - 4 = 8P_{p^a - (\frac{2}{p^a})}^2 \equiv 0 \pmod{p^2}$$

and hence

$$4 \left(\frac{2}{p^a}\right) P_{p^a} - Q_{p^a} = \left(\frac{2}{p^a}\right) Q_{p^a - (\frac{2}{p^a})} \equiv 2 \pmod{p^2}.$$

It follows that

$$\begin{aligned} & \left(\frac{2}{p^a}\right) P_{p^a - (\frac{2}{p^a})} Q_{p^a - (\frac{2}{p^a})} \\ & \equiv 2P_{p^a - (\frac{2}{p^a})} = Q_{p^a} - 2 \left(\frac{2}{p^a}\right) P_{p^a} \equiv \frac{Q_{p^a}}{2} - 1 \pmod{p^2}. \end{aligned}$$

This proves (2.1). $\square$

Lemma 2.2. *Let p be an odd prime and let a be a positive integer. Suppose that $p \equiv 1 \pmod{4}$ or $a > 1$. Then*

$$p^a \sum_{0 \leq k < \lfloor p^a/4 \rfloor} \frac{1}{\binom{(p^a-3)/2}{k}} \equiv \left(\frac{2}{p^a}\right) \frac{Q_{p^a} - 2}{4} \pmod{p^2}. \quad (2.2)$$

Proof. If $p^a \equiv 1 \pmod{4}$, then $(p^a - 3)/2$ is odd and hence

$$\sum_{k=0}^{(p^a-1)/4-1} \frac{1}{\binom{(p^a-3)/2}{k}} = \frac{1}{2} \sum_{k=0}^{(p^a-3)/2} \frac{1}{\binom{(p^a-3)/2}{k}}.$$

If $p^a \equiv 3 \pmod{4}$, then $a \in \{3, 5, \dots\}$ and

$$\sum_{k=0}^{(p^a-3)/4-1} \frac{1}{\binom{(p^a-3)/2}{k}} = \frac{1}{2} \sum_{k=0}^{(p^a-3)/2} \frac{1}{\binom{(p^a-3)/2}{k}} - \frac{1}{2} \cdot \frac{1}{\binom{(p^a-3)/2}{(p^a-3)/4}}.$$

In the case $p^a \equiv 3 \pmod{4}$, as the fractional parts of $(p^a - 3)/(2p)$ and $(p^a - 3)/(4p)$ are $(p - 3)/(2p)$ and $(p - 3)/(4p)$ respectively, we have

$$\left\lfloor \frac{(p^a - 3)/2}{p} \right\rfloor = 2 \left\lfloor \frac{(p^a - 3)/4}{p} \right\rfloor$$

and hence

$$\nu_p \left(\binom{(p^a - 3)/2}{(p^a - 3)/4} \right) = \sum_{j=1}^{a-1} \left(\left\lfloor \frac{(p^a - 3)/2}{p^j} \right\rfloor - 2 \left\lfloor \frac{(p^a - 3)/4}{p^j} \right\rfloor \right) < a - 1,$$

where $\nu_p(x)$ denotes the p -adic valuation of an integer x . (It is well known that $\nu_p(n!) = \sum_{j=1}^{\infty} \lfloor n/p^j \rfloor$ for any $n \in \mathbb{N}$.) Therefore,

$$p^a \sum_{0 \leq k < \lfloor p^a/4 \rfloor} \frac{1}{\binom{(p^a - 3)/2}{k}} \equiv \frac{p^a}{2} \sum_{k=0}^{(p^a - 3)/2} \frac{1}{\binom{(p^a - 3)/2}{k}} \pmod{p^2}.$$

Applying the known identity

$$\sum_{k=0}^n \frac{x^k}{\binom{n}{k}} = (n+1) \left(\frac{x}{1+x} \right)^{n+1} \sum_{k=1}^{n+1} \frac{1+x^k}{k(1+x)} \left(\frac{1+x}{x} \right)^k$$

(cf. [G, (2.4)]) with $x = 1$ and $n = (p^a - 3)/2$, we get

$$\begin{aligned} \frac{p^a}{2} \sum_{k=0}^{(p^a - 3)/2} \frac{1}{\binom{(p^a - 3)/2}{k}} &= \frac{p^a(p^a - 1)}{2^{(p^a + 3)/2}} \sum_{k=1}^{(p^a - 1)/2} \frac{2^k}{k} \\ &\equiv - \left(\frac{2}{p^a} \right) \frac{p^a}{4} \sum_{k=1}^{(p^a - 1)/2} \frac{2^k}{k} \pmod{p^2}. \end{aligned}$$

Since

$$\binom{p^a}{j} = \frac{p^a}{j} \prod_{0 < i < j} \frac{p^a - i}{i} \equiv \frac{p^a}{j} (-1)^{j-1} \pmod{p^2}$$

for all $j = 1, \dots, p^a$, we have

$$\begin{aligned} p^a \sum_{k=1}^{(p^a - 1)/2} \frac{2^k}{k} &\equiv -2 \sum_{k=1}^{(p^a - 1)/2} \binom{p^a}{2k} 2^k = - \sum_{k=1}^{p^a} \binom{p^a}{k} (\sqrt{2}^k + (-\sqrt{2})^k) \\ &= -(1 + \sqrt{2})^{p^a} - (1 - \sqrt{2})^{p^a} + 2 = -Q_{p^a} + 2 \pmod{p^2}. \end{aligned}$$

Combining the above we immediately get (2.2). $\square$

Proof of Theorem 1.1. Clearly,

$$\binom{-1/2}{k} = \frac{\binom{2k}{k}}{(-4)^k} \quad \text{for all } k = 0, 1, 2, \dots$$

Choose $\delta \in \{1, 3\}$ such that $p^a \equiv \delta \pmod{4}$. Then

$$\sum_{k=0}^{\lfloor \frac{3}{4}p^a \rfloor} \binom{-1/2}{k} = \sum_{k=0}^{p^a-1} \frac{\binom{2k}{k}}{(-4)^k} - \sum_{k=(3p^a+\delta)/4}^{p^a-1} \frac{\binom{2k}{k}}{(-4)^k}.$$

By Sun [S1, Theorem 1.1],

$$\sum_{k=0}^{p^a-1} \frac{\binom{2k}{k}}{(-4)^k} \equiv \left(\frac{2}{p^a} \right) + u_{p^a - (\frac{2}{p^a})}(-6, 1) \pmod{p^2}.$$

Hence we only need to prove the following congruence

$$\sum_{k=(3p^a+\delta)/4}^{p^a-1} \frac{\binom{2k}{k}}{(-4)^k} \equiv u_{p^a - (\frac{2}{p^a})}(-6, 1) \pmod{p^2} \quad (2.3)$$

provided that $p \equiv 1 \pmod{4}$ or $a > 1$.

Let k and l be positive integers with $k+l = p^a$ and $0 < l < p^a/2$. Then

$$\frac{\binom{2k}{k}}{\binom{2p^a-2}{p^a-1}} = \frac{(2p^a-2l)!}{(2p^a-2)!} \left(\frac{(p^a-1)!}{(p^a-l)!} \right)^2 = \frac{\prod_{0 < i < l} (p^a-i)^2}{\prod_{1 < j < 2l} (2p^a-j)}$$

and hence

$$\frac{\binom{2k}{k}}{\binom{2p^a-2}{p^a-1}} \cdot \frac{(2l-1)!}{(l-1)!^2} = \frac{\prod_{0 < i < l} (1-p^a/i)^2}{\prod_{1 < j < 2l} (1-2p^a/j)} \equiv 1 \pmod{p}.$$

Note that

$$\binom{2p^a-2}{p^a-1} = p^a \prod_{j=2}^{p^a-1} \frac{2p^a-j}{j} \equiv -p^a \pmod{p^{a+1}}$$

and

$$\binom{2k}{k} = \binom{p^a + (2k-p^a)}{0p^a+k} \equiv \binom{2k-p^a}{k} = 0 \pmod{p}$$

by Lucas' theorem. So we have

$$\frac{l}{2} \binom{2l}{l} = \frac{(2l-1)!}{(l-1)!^2} \not\equiv 0 \pmod{p^a}$$

and

$$\binom{2k}{k} \equiv -p^a \frac{(l-1)!^2}{(2l-1)!} = -\frac{2p^a}{l \binom{2l}{l}} \pmod{p^2}.$$

In view of the above,

$$\sum_{k=(3p^a+\delta)/4}^{p^a-1} \frac{\binom{2k}{k}}{(-4)^k} \equiv \frac{-2p^a}{(-4)^{p^a}} \sum_{l=1}^{(p^a-\delta)/4} \frac{(-4)^l}{l \binom{2l}{l}} \equiv \frac{p^a}{2} \sum_{k=1}^{(p^a-\delta)/4} \frac{(-4)^k}{k \binom{2k}{k}} \pmod{p^2}.$$

For $k = 1, \dots, (p^a - 1)/2$, clearly

$$\begin{aligned} \frac{\binom{(p^a-1)/2}{k}}{\binom{2k}{k}/(-4)^k} &= \frac{\binom{(p^a-1)/2}{k}}{\binom{-1/2}{k}} = \prod_{j=0}^{k-1} \frac{(p^a-1)/2-j}{-1/2-j} \\ &= \prod_{j=0}^{k-1} \left(1 - \frac{p^a}{2j+1}\right) \equiv 1 \pmod{p} \end{aligned}$$

and hence

$$\frac{\binom{(p^a-3)/2}{k-1}}{k \binom{2k}{k}/(-4)^k} \equiv \frac{2}{p^a-1} \equiv -2 \pmod{p}.$$

Therefore

$$\frac{p^a}{2} \sum_{k=1}^{(p^a-\delta)/4} \frac{(-4)^k}{k \binom{2k}{k}} \equiv -p^a \sum_{k=0}^{(p^a-\delta)/4-1} \frac{1}{\binom{(p^a-3)/2}{k}} \pmod{p^2}.$$

So far we have reduced (2.3) to the following congruence:

$$p^a \sum_{k=0}^{(p^a-\delta)/4-1} \frac{1}{\binom{(p^a-3)/2}{k}} \equiv -u_{p^a-(\frac{2}{p^a})}(-6, 1) \pmod{p^2}. \quad (2.4)$$

In view of (2.4) and Lemma 2.2, it suffices to show that

$$u_{p^a-(\frac{2}{p^a})}(-6, 1) \equiv -\left(\frac{2}{p^a}\right) \frac{Q_{p^a}-2}{4} \pmod{p^2}.$$

As $-3+2\sqrt{2}$ and $-3-2\sqrt{2}$ are the two roots of the equation $x^2+6x+1=0$, for any $n \in \mathbb{N}$ we have

$$\begin{aligned} u_n(-6, 1) &= \frac{(-3+2\sqrt{2})^n - (-3-2\sqrt{2})^n}{4\sqrt{2}} \\ &= \frac{(-1)^{n-1}}{2} \cdot \frac{(1+\sqrt{2})^{2n} - (1-\sqrt{2})^{2n}}{2\sqrt{2}} = \frac{(-1)^{n-1}}{2} P_n Q_n \end{aligned}$$

Therefore, with the help of Lemma 2.1 we finally obtain

$$u_{p^a-(\frac{2}{p^a})}(-6, 1) = -\frac{1}{2} P_{p^a-(\frac{2}{p^a})} Q_{p^a-(\frac{2}{p^a})} \equiv -\left(\frac{2}{p^a}\right) \frac{Q_{p^a}-2}{4} \pmod{p^2}$$

as desired.

The proof of Theorem 1.1 is now complete. $\square$

3. PROOF OF THEOREM 1.2

Lemma 3.1. *Let $p > 3$ be a prime. Then we have the following congruence:*

$$\left(\frac{x^p + (1-x)^p - 1}{p} \right)^2 \equiv -2 \sum_{k=1}^{p-1} \frac{(1-x)^k}{k^2} - 2x^{2p} \sum_{k=1}^{p-1} \frac{(1-x^{-1})^k}{k^2} \pmod{p}. \quad (3.1)$$

Proof. (3.1) follows immediately if we combine (4) and (5) of Granville [Gr]. $\square$

Proposition 3.1. *Let A and B be nonzero integers, and let α and β be the two roots of the equation $x^2 - Ax + B = 0$. Let p be an odd prime not dividing AB . Then*

$$\left(\frac{v_p(A, B) - A^p}{p} \right)^2 \equiv -2A^2 \sum_{k=1}^{p-1} \frac{\alpha^k}{A^k k^2} - 2\beta^{2p} \sum_{k=1}^{p-1} \frac{\alpha^{2k}}{(-B)^k k^2} \pmod{p}, \quad (3.2)$$

and

$$\left(\frac{v_p(A, B) - A^p}{p} \right)^2 \equiv -2A\alpha^p \sum_{k=1}^{p-1} \frac{\alpha^k}{A^k k^2} - 2\beta^{2p} \sum_{k=1}^{p-1} \frac{A^k \alpha^k}{B^k k^2} \pmod{p}. \quad (3.3)$$

Proof. By (3.1) and Fermat's little theorem,

$$\begin{aligned} & \frac{1}{A^2} \left(\frac{x^p + (A-x)^p - A^p}{p} \right)^2 \\ & \equiv \left(\frac{(x/A)^p + (1-x/A)^p - 1}{p} \right)^2 \\ & \equiv -2 \sum_{k=1}^{p-1} \frac{(1-x/A)^k}{k^2} - 2 \left(\frac{x}{A} \right)^{2p} \sum_{k=1}^{p-1} \frac{(1-A/x)^k}{k^2} \pmod{p}. \end{aligned}$$

Note that $v_p(A, B) = \beta^p + \alpha^p = \beta^p + (A - \beta)^p$ and $\alpha\beta = B$. So we have

$$\begin{aligned} \left(\frac{v_p(A, B) - A^p}{p} \right)^2 & \equiv -2A^2 \sum_{k=1}^{p-1} \frac{(A - \beta)^k}{A^k k^2} \\ & \quad - 2\beta^{2p} \sum_{k=1}^{p-1} \frac{(1 - A\alpha/B)^k}{k^2} \pmod{p} \end{aligned}$$

and hence (3.2) holds since $A\alpha - B = \alpha^2$.

On the other hand,

$$\alpha^p(A^p - v_p(A, B)) = \alpha^p(A^p - \alpha^p - \beta^p) = (B + \alpha^2)^p + (-\alpha^2)^p - B^p$$

and hence

$$\begin{aligned} & \alpha^{2p} \left(\frac{A^p - v_p(A, B)}{p} \right)^2 \\ &= \left(\frac{(-\alpha^2)^p + (B - (-\alpha^2))^p - B^p}{p} \right)^2 \\ &\equiv -2B^2 \sum_{k=1}^{p-1} \frac{(1 - (-\alpha^2)/B)^k}{k^2} - 2(-\alpha^2)^{2p} \sum_{k=1}^{p-1} \frac{(1 - B/(-\alpha^2))^k}{k^2} \\ &= -2B^2 \sum_{k=1}^{p-1} \frac{(A\alpha)^k}{B^k k^2} - 2\alpha^{4p} \sum_{k=1}^{p-1} \frac{(A\alpha)^k}{\alpha^{2k} k^2} \\ &\equiv -2(\alpha\beta)^{2p} \sum_{k=1}^{p-1} \frac{(A\alpha)^k}{B^k k^2} - 2A\alpha^{3p} \sum_{k=1}^{p-1} \frac{\alpha^{p-k}}{A^{p-k}(p-k)^2} \pmod{p}. \end{aligned}$$

Therefore (3.3) follows. $\square$

Proof of Theorem 1.2. Let α and β be the two roots of the equation $x^2 - x - 1 = 0$. Applying Proposition 3.1 with $A = 1$ and $B = -1$, we get

$$\left(\frac{L_p - 1}{p} \right)^2 \equiv -2 \sum_{k=1}^{p-1} \frac{\alpha^k}{k} - 2\beta^{2p} \sum_{k=1}^{p-1} \frac{\alpha^{2k}}{k^2} \pmod{p} \quad (3.4)$$

and

$$\left(\frac{L_p - 1}{p} \right)^2 \equiv -2\alpha^p \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} - 2\beta^{2p} \sum_{k=1}^{p-1} \frac{(-\alpha)^k}{k^2} \pmod{p}. \quad (3.5)$$

Since

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{2\alpha^{2k}}{(2k)^2} &= \sum_{j=1}^{2p-1} (1 + (-1)^j) \frac{\alpha^j}{j^2} = \sum_{k=1}^{p-1} \left(\frac{\alpha^k + (-\alpha)^k}{k^2} + \frac{\alpha^{p+k} + (-\alpha)^{p+k}}{(p+k)^2} \right) \\ &\equiv (1 + \alpha^p) \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} + (1 - \alpha^p) \sum_{k=1}^{p-1} \frac{(-\alpha)^k}{k^2} \pmod{p}, \end{aligned}$$

(3.4) can be rewritten as

$$\begin{aligned} \left(\frac{L_p - 1}{p} \right)^2 &\equiv -2(1 + 2(1 + \alpha^p)\beta^{2p}) \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} \\ &\quad - 4(1 - \alpha^p)\beta^{2p} \sum_{k=1}^{p-1} \frac{(-\alpha)^k}{k^2} \pmod{p}. \end{aligned} \quad (3.6)$$

Multiplying (3.5) by $2(1 - \alpha^p)$ and then subtracting it from (3.6) we obtain

$$\begin{aligned} (2\alpha^p - 1) \left(\frac{L_p - 1}{p} \right)^2 &\equiv (4\alpha^p(1 - \alpha^p) - 2 - 4(1 + \alpha^p)\beta^{2p}) \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} \\ &= (4L_p - 4L_{2p} - 2) \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} \pmod{p}. \end{aligned}$$

Now that $L_p \equiv 1 \pmod{p}$ and

$$L_{2p} = \alpha^{2p} + \beta^{2p} \equiv (\alpha^2 + \beta^2)^p = ((\alpha + \beta)^2 - 2\alpha\beta)^p = 3^p \equiv 3 \pmod{p},$$

we have

$$(2\alpha^p - 1) \left(\frac{L_p - 1}{p} \right)^2 \equiv (4 - 4 \times 3 - 2) \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} = -10 \sum_{k=1}^{p-1} \frac{\alpha^k}{k^2} \pmod{p}.$$

Similarly,

$$(2\beta^p - 1) \left(\frac{L_p - 1}{p} \right)^2 \equiv -10 \sum_{k=1}^{p-1} \frac{\beta^k}{k^2} \pmod{p}. \quad (3.7)$$

As $2\alpha^p - 1 + (2\beta^p - 1) = 2L_p - 2 \equiv 0 \pmod{p}$, we finally obtain

$$\sum_{k=1}^{p-1} \frac{L_k}{k^2} = \sum_{k=1}^{p-1} \frac{\alpha^k + \beta^k}{k^2} \equiv 0 \pmod{p}.$$

So far we have completed the proof of Theorem 1.2. $\square$

Remark 3.1. Let $p > 5$ be a prime. In view of (3.7), we also have

$$\sum_{k=1}^{p-1} \frac{F_k}{k^2} \equiv -\frac{2F_p}{10} \left(\frac{L_p - 1}{p} \right)^2 \equiv -\frac{1}{5} \left(\frac{p}{5} \right) \left(\frac{L_p - 1}{p} \right)^2 \pmod{p}. \quad (3.8)$$

4. PROOF OF THEOREM 1.3

We need a lemma which extends a congruence due to Granville [Gr, (6)].

Lemma 4.1. *Let p be an odd prime and let a be a positive integer. Then*

$$\begin{aligned} p\delta_{p,3} + p^{a-1} \sum_{k=1}^{p^a-1} \frac{(1-x)^k}{k} \\ \equiv \frac{1 - x^{p^a} - (1-x)^{p^a}}{p} - p \left(\sum_{k=1}^{p-1} \frac{x^k}{k^2} \right)^{p^{a-1}} \pmod{p^2}. \end{aligned} \quad (4.1)$$

Proof. As observed by Granville [Gr], for any integer $n > 1$ we have

$$\begin{aligned}
 \sum_{k=1}^{n-1} \frac{(1-x)^k - 1}{k} &= \sum_{k=1}^{n-1} \frac{1}{k} \sum_{j=1}^k \binom{k}{j} (-x)^j = \sum_{j=1}^{n-1} \frac{(-x)^j}{j} \sum_{k=j}^{n-1} \binom{k-1}{j-1} \\
 &= \sum_{j=1}^{n-1} \frac{(-x)^j}{j} \binom{n-1}{j} \quad (\text{by [G, (1.52)]}) \\
 &= \sum_{j=1}^{n-1} \frac{(-x)^j}{j} \left(\frac{n}{j} \binom{n-1}{j-1} - \frac{j}{n} \binom{n}{j} \right) \\
 &= n \sum_{j=1}^{n-1} \binom{n-1}{j-1} \frac{(-x)^j}{j^2} - \frac{(1-x)^n - (-x)^n - 1}{n}.
 \end{aligned}$$

Note that

$$\begin{aligned}
 2p^{a-1} \sum_{k=1}^{p^a-1} \frac{1}{k} &= p^{a-1} \sum_{k=1}^{p^a-1} \left(\frac{1}{k} + \frac{1}{p^a - k} \right) = \sum_{k=1}^{p^a-1} \frac{p^a}{k} \cdot \frac{p^{a-1}}{p^a - k} \\
 &\equiv \sum_{j=1}^{p-1} \frac{p^a}{p^{a-1}j} \cdot \frac{p^{a-1}}{p^a - p^{a-1}j} \equiv -p \sum_{j=1}^{p-1} \frac{1}{j^2} \equiv p\delta_{p,3} \pmod{p^2}.
 \end{aligned}$$

(As $\sum_{j=1}^{p-1} 1/j^2 \equiv \sum_{k=1}^{p-1} 1/(2k)^2 \pmod{p}$, we have $\sum_{j=1}^{p-1} 1/j^2 \equiv 0 \pmod{p}$ if $p > 3$.) Hence

$$p^{a-1} \sum_{k=1}^{p^a-1} \frac{1}{k} \equiv -p\delta_{p,3} \pmod{p^2}.$$

Note also that

$$\binom{p^a-1}{j-1} = \prod_{0 < i < j} \frac{p^a - i}{i} \equiv (-1)^{j-1} \pmod{p}.$$

Therefore

$$\begin{aligned}
 &p^{a-1} \sum_{k=1}^{p^a-1} \frac{(1-x)^k}{k} + p\delta_{p,3} \\
 &\equiv p^{a-1} \sum_{k=1}^{p^a-1} \frac{(1-x)^k - 1}{k} \\
 &\equiv -p^{2a-1} \sum_{j=1}^{p^a-1} \frac{x^j}{j^2} - \frac{x^{p^a} + (1-x)^{p^a} - 1}{p} \pmod{p^2}
 \end{aligned}$$

To complete the proof, it suffices to note that

$$p^{2a-1} \sum_{j=1}^{p^a-1} \frac{x^j}{j} \equiv p^{2a-1} \sum_{k=1}^{p-1} \frac{x^{p^{a-1}k}}{(p^{a-1}k)^2} = p \sum_{k=1}^{p-1} \frac{x^{p^{a-1}k}}{k^2} \pmod{p^2}$$

and

$$\left(\sum_{k=1}^{p-1} \frac{x^k}{k^2} \right)^{p^{a-1}} \equiv \sum_{k=1}^{p-1} \left(\frac{x^k}{k^2} \right)^{p^{a-1}} \equiv \sum_{k=1}^{p-1} \frac{x^{p^{a-1}k}}{k^2} \pmod{p}.$$

We are done. $\square$

Proposition 4.1. *Let $p \neq 2, 5$ be a prime and let a be a positive integer. Then*

$$p^{a-1} \sum_{k=1}^{p^a-1} \frac{F_{2(p^a-k)}}{k} \equiv \frac{F_{2p^a} - F_{p^a}}{p} + \frac{p}{10} \left(\frac{p^a}{5} \right) \left(\frac{L_p - 1}{p} \right)^2 \pmod{p^2}. \quad (4.2)$$

Proof. Let α and β be the two roots of the equation $x^2 - x - 1 = 0$. Clearly $\alpha + \beta = 1$ and $\alpha\beta = -1$. Set

$$g(x) = p^{a-1} \sum_{k=1}^{p^a-1} \frac{x^k}{k}, \quad q(x) = \frac{x^{p^a} + (1-x)^{p^a} - 1}{p} \quad \text{and} \quad G(x) = \sum_{k=1}^{p-1} \frac{x^k}{k^2}.$$

By Lemma 4.1 we have

$$p\delta_{p,3} + g(1-x) \equiv -q(x) - pG(x)^{p^{a-1}} \pmod{p^2}.$$

In view of (3.7),

$$G(\beta) \equiv \frac{1-2\beta^p}{10} \left(\frac{L_p-1}{p} \right)^2 - \delta_{p,3} \pmod{p}$$

(this can be checked directly when $p = 3$). Hence

$$\begin{aligned} G(-\alpha) &= G(\beta^{-1}) = \frac{1}{\beta^p} \sum_{k=1}^{p-1} \frac{\beta^{p-k}}{k^2} \\ &\equiv -\alpha^p G(\beta) \equiv \delta_{p,3} \alpha^p - \frac{2+\alpha^p}{10} \left(\frac{L_p-1}{p} \right)^2 \pmod{p}. \end{aligned}$$

Note that

$$q(-\alpha) = \frac{(-\alpha)^{p^a} + (1+\alpha)^{p^a} - 1}{p} = \frac{\alpha^{p^a}(\alpha^{p^a} + \beta^{p^a} - 1)}{p} = \alpha^{p^a} \frac{L_{p^a} - 1}{p}.$$

Applying Lemma 4.1 we get

$$\begin{aligned}
 g(\alpha^2) &= g(1 - (-\alpha)) \\
 &\equiv -p\delta_{p,3} - pG(-\alpha)^{p^{a-1}} - q(-\alpha) \\
 &\equiv -p\delta_{p,3} - p\delta_{p,3}\alpha^{p^a} + p\left(\frac{\alpha^p + 2}{10}\right)^{p^{a-1}} \left(\frac{L_p - 1}{p}\right)^{2p^{a-1}} - \alpha^{p^a} \frac{L_{p^a} - 1}{p} \\
 &\equiv -p\delta_{p,3}(1 + \alpha)^{p^a} + p\frac{\alpha^{p^a} + 2}{10} \left(\frac{L_p - 1}{p}\right)^2 - \alpha^{p^a} \frac{L_{p^a} - 1}{p} \pmod{p^2}
 \end{aligned}$$

and hence

$$\begin{aligned}
 &\beta^{2p^a} g(\alpha^2) + p\delta_{p,3}(\alpha\beta)^{2p^a} \\
 &\equiv p\frac{2\beta^{2p^a} - \beta^{p^a}}{10} \left(\frac{L_p - 1}{p}\right)^2 + \beta^{p^a} \frac{L_{p^a} - 1}{p} \pmod{p^2}.
 \end{aligned}$$

As $\beta^{2p^a} = (1 + \beta)^{p^a} \equiv 1 + \beta^{p^a} \pmod{p}$, we have

$$\beta^{2p^a} g(\alpha^2) \equiv p\delta_{p,3} + p\frac{2 + \beta^{p^a}}{10} \left(\frac{L_p - 1}{p}\right)^2 + \beta^{p^a} \frac{L_{p^a} - 1}{p} \pmod{p^2}.$$

Similarly,

$$\alpha^{2p^a} g(\beta^2) \equiv p\delta_{p,3} + p\frac{2 + \alpha^{p^a}}{10} \left(\frac{L_p - 1}{p}\right)^2 + \alpha^{p^a} \frac{L_{p^a} - 1}{p} \pmod{p^2}.$$

Observe that

$$\begin{aligned}
 \sum_{k=1}^{p^a-1} \frac{F_{2(p^a-k)}}{k} &= \sum_{k=1}^{p^a-1} \frac{\alpha^{2p^a-2k} - \beta^{2p^a-2k}}{(\alpha - \beta)k} \\
 &= \frac{1}{\alpha - \beta} \left(\alpha^{2p^a} \sum_{k=1}^{p^a-1} \frac{\beta^{2k}}{k} - \beta^{2p^a} \sum_{k=1}^{p^a-1} \frac{\alpha^{2k}}{k} \right).
 \end{aligned}$$

So, by the above, we have

$$\begin{aligned}
 p^{a-1} \sum_{k=1}^{p^a-1} \frac{F_{2(p^a-k)}}{k} &= \frac{\alpha^{2p^a} g(\beta^2) - \beta^{2p^a} g(\alpha^2)}{\alpha - \beta} \\
 &\equiv p\frac{F_{p^a}}{10} \left(\frac{L_p - 1}{p}\right)^2 + F_{p^a} \frac{L_{p^a} - 1}{p} \\
 &\equiv \frac{p}{10} \left(\frac{p^a}{5}\right) \left(\frac{L_p - 1}{p}\right)^2 + \frac{F_{2p^a} - F_{p^a}}{p} \pmod{p^2}.
 \end{aligned}$$

This concludes the proof. $\square$

Lemma 4.2. *Let $p \neq 2, 5$ be a prime and let a be a positive integer. Then*

$$\left(\frac{p^a}{5}\right) (2F_{p^a} - F_{2p^a}) + \frac{(L_p - 1)^2}{5} \equiv 1 - 2F_{p^a - (\frac{p^a}{5})} \pmod{p^3}. \quad (4.3)$$

Proof. Note that

$$(L_{p^a} - 1)^2 = p^2 \left(\frac{L_{p^a} - 1}{p}\right)^2 \equiv p^2 \left(\frac{L_p - 1}{p}\right)^2 = (L_p - 1)^2 \pmod{p^3}$$

since $L_{p^a} \equiv L_p \pmod{p^2}$ by [S1, (2.4)]. Also,

$$L_{p^a} = F_{p^a} + 2F_{p^a-1} = 2F_{p^a+1} - F_{p^a} = 2F_{p^a - (\frac{p^a}{5})} + \left(\frac{p^a}{5}\right) F_{p^a}.$$

Thus

$$\begin{aligned} & 1 - 2F_{p^a - (\frac{p^a}{5})} - \left(\frac{p^a}{5}\right) (2F_{p^a} - F_{2p^a}) \\ &= 1 - L_{p^a} + \left(\frac{p^a}{5}\right) F_{p^a} - \left(\frac{p^a}{5}\right) (2F_{p^a} - F_{p^a} L_{p^a}) \\ &= (L_{p^a} - 1) \left(\left(\frac{p^a}{5}\right) F_{p^a} - 1\right) \end{aligned}$$

and hence it suffices to show that

$$\left(\frac{p^a}{5}\right) F_{p^a} - 1 \equiv \frac{L_{p^a} - 1}{5} \pmod{p^2} \quad (4.4)$$

as $L_{p^a} \equiv 1 \pmod{p}$. Since $L_{p^a} \equiv L_p \pmod{p^2}$, and

$$F_{p^a} \equiv \left(\frac{p}{5}\right)^{a-1} F_p \pmod{p^2}$$

by [S1, (2.5)], (4.4) is reduced to the case $a = 1$. Note that

$$\left(\frac{p}{5}\right) F_p - 1 = L_p - 2F_{p - (\frac{p}{5})} - 1 \equiv \frac{L_p - 1}{5} \pmod{p^2}$$

since $L_p \equiv 1 + \frac{2}{5}F_{p - (\frac{p}{5})} \pmod{p^2}$ by the proof of [ST, Corollary 1.3]. We are done. $\square$

Proof of Theorem 1.3. Applying [ST, (2.4)] with $d = 0$ and $n = p^a$, we get

$$\sum_{k=0}^{p^a-1} (-1)^k \binom{2k}{k} = \sum_{k=0}^{p^a-1} (-1)^k \binom{2p^a}{k} F_{2(p^a-k)}. \quad (4.5)$$

For each $k = 1, \dots, p^a - 1$, clearly

$$\begin{aligned} (-1)^k \binom{2p^a}{k} &= (-1)^k \frac{2p^a}{k} \binom{2p^a - 1}{k - 1} = -\frac{2p^a}{k} \prod_{0 < j < k} \left(1 - \frac{2p^a}{j}\right) \\ &\equiv -\frac{2p^a}{k} \left(1 - 2p^a \sum_{0 < j < k} \frac{1}{j}\right) \pmod{p^3} \end{aligned}$$

and similarly

$$(-1)^k \binom{p^a}{k} \equiv -\frac{p^a}{k} \left(1 - p^a \sum_{0 < j < k} \frac{1}{j}\right) \pmod{p^3},$$

hence

$$(-1)^k \binom{2p^a}{k} \equiv 4(-1)^k \binom{p^a}{k} + \frac{2p^a}{k} \pmod{p^3}.$$

So, (4.5) yields that

$$\begin{aligned} &\sum_{k=0}^{p^a-1} (-1)^k \binom{2k}{k} - F_{2p^a} \\ &\equiv 4 \sum_{k=1}^{p^a-1} \binom{p^a}{k} (-1)^k F_{2(p^a-k)} + 2 \sum_{k=1}^{p^a-1} \frac{p^a}{k} F_{2(p^a-k)} \\ &= 4 \sum_{k=1}^{p^a-1} \binom{p^a}{k} (-1)^{p^a-k} F_{2k} + 2 \sum_{k=1}^{p^a-1} \frac{p^a}{k} F_{2(p^a-k)} \pmod{p^3}. \end{aligned}$$

Let α and β be the two roots of the equation $x^2 - x - 1 = 0$. Then

$$\begin{aligned} \sum_{k=1}^{p^a-1} \binom{p^a}{k} (-1)^{p^a-k} F_{2k} &= -F_{2p^a} + \sum_{k=0}^{p^a} \binom{p^a}{k} (-1)^{p^a-k} \frac{\alpha^{2k} - \beta^{2k}}{\alpha - \beta} \\ &= \frac{(\alpha^2 - 1)^{p^a} - (\beta^2 - 1)^{p^a}}{\alpha - \beta} - F_{2p^a} = F_{p^a} - F_{2p^a}. \end{aligned}$$

Thus, by the above and Proposition 4.1 we have

$$\begin{aligned} \sum_{k=0}^{p^a-1} (-1)^k \binom{2k}{k} &\equiv F_{2p^a} + 4(F_{p^a} - F_{2p^a}) + 2p^a \sum_{k=1}^{p^a-1} \frac{F_{2(p^a-k)}}{k} \\ &\equiv 4F_{p^a} - 3F_{2p^a} + 2(F_{2p^a} - F_{p^a}) + \left(\frac{p^a}{5}\right) \frac{(L_p - 1)^2}{5} \\ &= 2F_{p^a} - F_{2p^a} + \left(\frac{p^a}{5}\right) \frac{(L_p - 1)^2}{5} \pmod{p^3}. \end{aligned}$$

Combining this with (4.3) we immediately get the desired (1.3). This completes the proof of Theorem 1.3. $\square$

Acknowledgment. After we established Theorem 1.2, we got Roberto Tauraso's generous permission (on Oct. 1, 2010) to cite his conjecture (1.2) (cf. [T]). We would like to express our sincere thanks to Prof. R. Tauraso.

REFERENCES

- [CDP] R. Crandall, K. Dilcher and C. Pomerance, *A search for Wieferich and Wilson primes*, Math. Comp. **66** (1997), 433–449.
- [CP] R. Crandall and C. Pomerance, *Prime Numbers: A Computational Perspective*, Second Edition, Springer, New York, 2005.
- [G] H. W. Gould, *Combinatorial Identities*, Morgantown Printing and Binding Co., 1972.
- [Gr] A. Granville, *The square of the Fermat quotient*, Integers **4** (2004), #A22, 3pp (electronic).
- [MR] R. J. McIntosh and E. L. Roettger, *A search for Fibonacci-Wieferich and Wolstenholme primes*, Math. Comp. **76** (2007), 2087–2094.
- [SS] Z. H. Sun and Z. W. Sun, *Fibonacci numbers and Fermat's last theorem*, Acta Arith. **60** (1992), 371–388.
- [S1] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients*, Sci. China Math. **53** (2010), 2473–2488. <http://arxiv.org/abs/0909.5648>.
- [S2] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients (II)*, preprint, arXiv:0911.3060. <http://arxiv.org/abs/0911.3060>.
- [ST] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45** (2010), 125–148.
- [T] R. Tauraso, *A personal communication via e-mail*, Jan. 29, 2010.